

CO₂ emissions from forest loss

G. R. van der Werf, D. C. Morton, R. S. DeFries, J. G. J. Olivier, P. S. Kasibhatla, R. B. Jackson, G. J. Collatz and J. T. Randerson

Deforestation is the second largest anthropogenic source of carbon dioxide to the atmosphere, after fossil fuel combustion. Following a budget reanalysis, the contribution from deforestation is revised downwards, but tropical peatlands emerge as a notable carbon dioxide source.

Programmes that aim to reduce the emissions from deforestation and forest degradation are being considered as a cost-effective way to mitigate anthropogenic greenhouse-gas emissions¹. Deforestation and forest degradation contribute to atmospheric greenhouse-gas emissions through combustion of forest biomass and decomposition of remaining plant material and soil carbon. Within the science and policy communities, carbon emissions from deforestation and forest degradation have been estimated to account for about 20% of global anthropogenic CO₂ emissions^{2–5}. A recalculation of this fraction using the same methods, but updated estimates on carbon emissions from both deforestation and fossil fuel combustion suggests that in 2008, the relative contribution of CO₂ emissions from deforestation and forest degradation was substantially smaller, around 12%. As a consequence, the maximum carbon savings from reductions in forest decline are likely to be lower than expected.

Deforestation is the long-term reduction of tree canopy cover⁶ to below 10–30%. In practice, deforestation is associated with the conversion of forest to other types of land use, such as cropland or pasture. Forest degradation is typically considered partial deforestation, with more than 10–30% of forest cover remaining (for example, through selective logging). Land degradation that does not involve changes in tree cover density, such as oxidation and combustion of deforested and drained tropical peatlands^{7,8}, may also involve substantial carbon emissions. However, losses of these non-forest carbon stocks are not generally included in deforestation and forest degradation assessments, as summarized in the 2007 report of the Intergovernmental Panel on Climate Change (IPCC) Working Group I (ref. 2), nor are they considered in policies aiming to reduce emissions from deforestation and forest degradation (REDD).

The two main assessments of CO₂ emissions from deforestation and




Figure 1 | Carbon emissions from deforestation and forest degradation (D&FD) and fossil fuel emissions for 1980 onwards. Updated datasets and approaches are depicted with a solid line, outdated ones with a dashed line. The often-held assumption that deforestation and forest degradation accounts for 20% of global anthropogenic CO₂ emissions is calculated using the average (dashed black line) of the deforestation surveys from the Food and Agricultural Organization (FAO; dashed grey line) and satellites in the 1990s (dark blue line), compared with average fossil fuel emissions estimates for the 1990s (red). Revised FAO-based emissions estimates (solid grey line) together with increased fossil fuel emissions substantially lower the relative contribution of deforestation and degradation to total anthropogenic emissions to about 12%. Dashed brown line indicates D&FD assessment used in IPCC Working Group III; solid brown line follows the same approach, but is based on updated carbon emissions from tropical forest fires used in this approach. Error bars are not shown, but the uncertainty in the deforestation figures is large, up to 50%. Peatland carbon emissions (0.30 Pg C yr⁻¹) are not included in emissions estimates.

forest degradation for the 1980s and 1990s — one based on surveys by the Food and Agricultural Organisation (FAO)⁹ and the other based on satellite data¹⁰ — have limitations^{11,12}, and one is not obviously preferable to the other. Therefore, IPCC Working Group I averaged emissions estimates from the two approaches for their estimate of carbon emissions from deforestation and forest

degradation (Fig. 1). However, since the IPCC assessments were written, the FAO has lowered its estimate of deforestation and forest degradation rates, leading to a 30% reduction in related carbon emissions^{13,14} (Fig. 1). Furthermore, an updated satellite-based estimate of deforestation rates¹⁵, derived from changes in tree cover density in the humid tropics during 2000–2005, suggested that rates

are similar to those in the previous decade¹⁰, resulting in estimates that are still lower than those reported in the revised FAO survey.

A recalculation, using IPCC methods and these updated values, suggests CO₂ emissions from deforestation and forest degradation (excluding peatland emissions) of about 1.2 Pg C yr⁻¹ — 23% less than the value given in the 2007 report of the IPCC Working Group I. In this calculation, we assume that rates of forest decline for 2000–2005 were almost the same as in the 1990s, based on survey¹⁴ and satellite-based approaches¹⁵ of deforestation area (Fig. 1). Carbon emissions from fossil fuel combustion have increased substantially over the same period^{16,17}, making the relative contribution from deforestation and forest degradation even smaller. As a result, deforestation and forest degradation emissions contribute about 12% of total anthropogenic CO₂ emissions (updated from 20%). Taking into account the large uncertainties in the deforestation and degradation estimates², the range becomes 6–17%.

In addition to the emissions assessment of the IPCC Working Group I (*The Physical Science Basis*), an independent assessment^{18,19} was carried out by the IPCC Working Group III (*Mitigation of Climate Change*), based on carbon emissions released by fires in tropical forest areas. This approach assumed that half of the total CO₂ emissions from deforestation and forest degradation come from forest fire, whereas the remainder can be attributed to respiration of leftover materials^{18,20}. Carbon emissions from tropical peatland burning and oxidation in Southeast Asia⁸, mostly in Indonesia, were included in their assessment. Combined with deforestation and forest degradation, this estimate of the contribution of forest and peatland decline amounted to 23% of total CO₂ emissions (or 17% of all anthropogenic greenhouse gases released into the atmosphere; see Supplementary Information).

However, the IPCC Working Group III assessments also need to be revised down in the light of new information: data updates led to a 32% decrease in the estimates of fire-related carbon emissions from deforestation and forest degradation, resulting in a revised deforestation and forest degradation average of 1.2 Pg C yr⁻¹ over the period 1997–2006 (refs 17, 21). Furthermore, carbon emissions from Southeast-Asian peatland fires, constrained by satellite-derived column-mixing ratios of carbon monoxide²², were less than half of those used in the IPCC Working Group III report. Yet even this lower value of the

contribution from peat fires highlights the importance of peat degradation in Southeast Asia in the global CO₂ budget: combined with earlier estimates of carbon emissions from peatland oxidation⁸, it amounts to 0.30 Pg C yr⁻¹ averaged over 1997–2006, about a quarter of the updated CO₂ emissions from deforestation and forest degradation. The combined contribution of deforestation, forest degradation and peatland emissions to total anthropogenic CO₂ emissions is about 15% (range 8–20%; see Supplementary Information for a comparison with all anthropogenic greenhouse gases). We conclude that taking into account carbon emissions from peatlands would enhance the effectiveness of REDD programmes, which are under discussion in the United Nation's climate policy negotiations.

In short, the maximum reduction in CO₂ emissions from avoiding deforestation and forest degradation is probably about 12% of current total anthropogenic emissions (or 15% if peat degradation is included) — and that is assuming, unrealistically, that emissions from deforestation, forest degradation and peat degradation can be completely eliminated. Therefore, reducing fossil fuel emissions remains the key element for stabilizing atmospheric CO₂ concentrations.

Nevertheless, efforts to mitigate emissions from tropical forests and peatlands, and maintain existing terrestrial carbon stocks, remain critical for the negotiation of a post-Kyoto agreement. Even our revised estimates represent substantial emissions, and for about 30 developing countries, including Brazil, Bolivia, Indonesia, Myanmar and Zambia, deforestation and forest degradation are the largest source of CO₂ (ref. 17). Moreover, reductions in the emissions from deforestation and degradation of peat and forest may remain one of the more cost-effective 'wedges'^{11,23} that can help to stabilize atmospheric CO₂ levels.

If changes in terrestrial carbon storage are to have a role in a post-Kyoto agreement, a strong focus on monitoring changes in carbon content, irrespective of forest cover density would strengthen the effectiveness of REDD programmes. For example, replacing peat forest with oil-palm plantations may not change the tree cover density, but it does lead to a large pulse of CO₂ emissions because of reductions in both tree biomass and soil carbon⁸. In addition to the proposed satellite monitoring of terrestrial biomass, satellite-derived observations of fires as well as measurements of atmospheric CO₂ concentrations in combination with inverse modelling and ground-based validation can

help track carbon losses from deforestation and the degradation of forest and peat. □

G. R. van der Werf^{1*}, D. C. Morton², R. S. DeFries³, J. G. J. Olivier⁴, P. S. Kasibhatla⁵, R. B. Jackson⁵, G. J. Collatz², and J. T. Randerson⁶ are at the ¹Faculty of Earth and Life Sciences, VU University Amsterdam, De Boelelaan 1085, 1081HV, Amsterdam, The Netherlands, ²Hydrospheric and Biospheric Sciences Laboratory, Code 614.4, NASA's Goddard Space Flight Center, Greenbelt, Maryland 20771, USA, ³Department of Ecology, Evolution and Environmental Biology, Columbia University, 10th Floor Schermerhorn Extension, 1200 Amsterdam Avenue, New York, New York 10027, USA, ⁴Netherlands Environmental Assessment Agency, PO Box 303, 3720AH Bilthoven, The Netherlands, ⁵Nicholas School of the Environment, Duke University, Durham, North Carolina 27708, USA and ⁶Earth System Science Department, 3212 Croul Hall, University of California, Irvine, California 92697, USA.

*e-mail: guido.van.der.werf@falw.vu.nl

References

- Gullison, R. E. *et al.* *Science* **316**, 985–986 (2007).
- Denman, K. L. *et al.* in *IPCC Climate Change 2007: The Physical Science Basis* (eds Solomon, S. *et al.*) 499–587 (Cambridge Univ. Press, 2007).
- Gibbs, H. K. & Herold, M. *Env. Res. Lett.* **2**, 045021 (2007).
- Schrope, M. *Nat. Rep. Clim. Change* **3**, 101–103 (2009).
- Available via <<http://tinyurl.com/kszq9t>>.
- Penman, J. *et al.* (eds) *Good Practice Guide for Land Use, Land-Use Change and Forestry* (Institute for Global Environmental Strategies, 2003).
- Page, S. E. *et al.* *Nature* **420**, 61–65 (2002).
- Hooijer, A. *et al.* *Delft Hydraulics report Q3943* (WL/Delft Hydraulics, 2006).
- Houghton, R. A. *Tellus B* **55**, 378–390 (2003).
- DeFries, R. S. *et al.* *Proc. Natl Acad. Sci. USA* **99**, 14256–14261 (2002).
- Grainger, A. *Proc. Natl Acad. Sci. USA* **105**, 818–823 (2008).
- Fearnside, P. M. & Laurance, W. F. *Science* **299**, 1015 (2003).
- Canadell, J. G. *et al.* *Proc. Natl Acad. Sci. USA* **104**, 18866–18870 (2007).
- Houghton, R. A. in *TRENDS: A Compendium of Data on Global Change* (Oak Ridge National Laboratory, US Department of Energy, 2008); available at <<http://cdiac.ornl.gov/trends/landuse/houghton/houghton.html>>.
- Hansen, M. C. *et al.* *Proc. Natl Acad. Sci. USA* **105**, 9439–9444 (2008).
- Andres, R. J. *et al.* *Tellus B* **51**, 759–765 (1999).
- Joint Research Centre/Netherlands Environmental Assessment Agency, *Emissions Database for Global Atmospheric Research 4.0* (2009); available at <<http://edgar.jrc.ec.europa.eu>>.
- Barker, T. *et al.* in *IPCC Climate Change 2007: Mitigation of Climate Change* (eds Metz, B. *et al.*) 25–93 (Cambridge Univ. Press, 2007).
- Nabuurs, G. J. *et al.* in *IPCC Climate Change 2007: Mitigation of Climate Change* (eds Metz, B. *et al.*) 541–584 (Cambridge Univ. Press, 2007).
- Olivier, J. G. J. *et al.* *Environ. Sci.* **2**, 81–99 (2005).
- van der Werf, G. R. *et al.* *Atmos. Chem. Phys.* **6**, 3423–3441 (2006).
- van der Werf, G. R. *et al.* *Proc. Natl Acad. Sci. USA* **105**, 20350–20355 (2008).
- Pacala, S. & Socolow, R. *Science* **305**, 968–972 (2004).
- Achard, F. *et al.* *Glob. Biogeochem. Cycles* **18**, GB2008 (2004).

Additional information

Supplementary information accompanies this paper on www.nature.com/geoscience.